

GOLDEN-THOMPSON FROM DAVIS

IGOR RIVIN

ABSTRACT. We give a very short proof of the Golden-Thompson inequality

The Golden-Thompson inequality (due independently to Golden [2] and Thompson [4]) states that for two hermitian matrices A and B , the following inequality holds:

$$(1) \quad \operatorname{tr} \exp(A + B) \leq \operatorname{tr} \exp A \operatorname{tr} \exp B.$$

The Golden-Thompson inequality (which is widely used in statistical physics and the theory of random matrices) is usually proved in a rather painful way using the power series of the matrix exponential function. In this note we point out that the Golden-Thompson follows immediately from the celebrated (in some circles) and underappreciated (in other circles) theorem of Chandler Davis, which states:

Theorem 1 (Chandler Davis' Theorem). *Let f be a unitarily invariant function on the set of hermitian matrices. Then f is convex if and only if its restriction to the set of diagonal matrices is a convex symmetric function.*

A function is *unitarily invariant* if $f(U^t A U) = f(A)$ for any unitary U . The original proof ([1]) is very short and elegant, and the reader is advised to read Davis' (one page) paper. A different two-page proof has been given by the author ([3]) Both proofs go through verbatim if we replace "hermitian" by "symmetric" and "unitary" by "orthogonal".

Now, the Golden-Thompson inequality states that the function $F(x) = \log \operatorname{tr} \exp(x)$ is convex on the set of hermitian matrices, and since it is obviously unitarily invariant, we need only check convexity of the function $f(\lambda_1, \dots, \lambda_n) = \log(\sum_{i=1}^n \exp(\lambda_i))$.

Date: October 12, 2010.

1991 Mathematics Subject Classification. 15A42, 15A52.

Key words and phrases. trace inequality, convexity.

Igor Rivin was supported by the National Science Foundation. He would also like to thank the Institute for Advanced Study for its hospitality and Nikhil Srivastava for interesting conversations.

Now, for the function f we can compute the hessian without too much trouble:

$$(2) \quad H_{ii} = \frac{\partial^2 f}{\partial \lambda_i^2} = \frac{\sum_{j \neq i} \exp(x_i + x_j)}{(\sum_{k=1}^n \exp(x_k))^2},$$

$$(3) \quad H_{ij} = \frac{\partial^2 f}{\partial \lambda_i \partial \lambda_j} = -\frac{\exp(x_i + x_j)}{(\sum_{k=1}^n \exp(x_k))^2}$$

It is easy to see that $H = DKD$, where D is the diagonal matrix with the i -th entry equal to $\exp(x_i)/(\sum_j \exp(x_j))$, and K is the Laplacian matrix of the complete graph on n vertices (in other words, $K_{ii} = n - 1$, and $K_{ij} = -1$ when $i \neq j$.) Since K is a connected graph, its laplacian matrix is positive semidefinite (with dimension of the null-space equal to 1), the Golden-Thompson inequality follows immediately.

REFERENCES

- [1] Chandler Davis. All convex invariant functions of hermitian matrices. *Arch. Math.*, 8:276–278, 1957.
- [2] Sidney Golden. Lower bounds for the Helmholtz function. *Phys. Rev. (2)*, 137:B1127–B1128, 1965.
- [3] Igor Rivin. Another simple proof of a theorem of Chandler Davis. *arXiv*, math.FA, Aug 2002. 3 pages.
- [4] Colin J. Thompson. Inequality with applications in statistical mechanics. *J. Mathematical Phys.*, 6:1812–1813, 1965.

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON
E-mail address: rivin@ias.edu